

Short communication

Modelling respiration of vegetation: evidence for a general temperature-dependent Q_{10}

MARK G. TJOELKER*, JACEK OLEKSYN^{†‡} and PETER B. REICH[†]

*Texas A & M University, Department of Forest Science, 2135 TAMU, College Station, TX 77843-2135, USA, †University of Minnesota, Department of Forest Resources, 1530 Cleveland Ave. N., St. Paul, MN 55108, USA, ‡Polish Academy of Sciences, Institute of Dendrology, Parkowa 5, PL-62-035 Kórnik, Poland

Abstract

Temperature responses of rates of respiratory CO₂ efflux from plants, soils, and ecosystems are frequently modelled using exponential functions with a constant Q_{10} near 2.0 (fractional change in rate with a 10 °C increase in temperature). However, we present evidence that Q_{10} declines with short-term increases in temperature in a predictable manner across diverse plant taxa. Thus, models using a constant Q_{10} are biased, and use of a temperature-corrected Q_{10} may improve the accuracy of modelled respiratory CO₂ efflux in plants and ecosystems in response to temperature and predicted global climate changes.

Keywords: biome, boreal tree species, dark respiration, elevated carbon dioxide, Q_{10} , temperature response function

Received 8 January 2000; revised version received 16 June 2000 and accepted 25 July 2000

Introduction

Biologists have long used Q_{10} to describe the temperature dependence of rates of biological processes, a concept originating in the 19th century physical-chemistry models of Arrhenius (1889) and Van't Hoff (1896). The Q_{10} function assumes an exponential relationship with temperature in which Q_{10} is the ratio of the rate at one temperature to that at a temperature 10 °C lower. Yet, it has long been known that short-term temperature responses of many biological processes, including respiration of plants (Belehrádek 1930; Wager 1941) and soils (Lloyd & Taylor 1994), often fail to accurately fit an exponential Q_{10} model over a broad range of temperatures. In plants, dark respiration (see reviews by James 1953; Forward 1960; Berry & Raison 1981) and respiratory enzymes (ap Rees *et al.* 1988; Farrar & Williams 1991) often exhibit declining values of Q_{10} with increasing measurement-temperature intervals. However, it remains unclear whether the temperature dependence of respiratory Q_{10} is a general phenomenon. The evidence suggests that respiratory Q_{10} is likely not constant, but

dependent on both the shape of the temperature-response curve and range of measurement temperatures used in its determination.

Despite these and other limitations, a simple exponential function (see equation in Materials and methods section) using a fixed Q_{10} of 2.0 has gained wide acceptance in modelling leaf- to ecosystem-scale respiration responses to temperature and global vegetation responses to climate change (Ryan 1991; Aber & Federer 1992; Melillo *et al.* 1993; Schimel *et al.* 1997; Cramer *et al.* 1999). Respiratory Q_{10} values are often reported near 2.0 (e.g. Amthor 1984; Collier & Cummins 1990). In a literature review, Q_{10} (10–20 °C) averaged 2.3 among 125 alpine, lowland, and crop species (Larigauderie & Körner 1995). However, few studies report Q_{10} values over a range of environmentally relevant temperatures for modelling temperature responses in contrasting climatic conditions.

Herein an experiment on the temperature dependence of respiratory Q_{10} for five boreal tree species is reported. In order to test further the temperature dependence of respiratory Q_{10} , literature data on the temperature responses of dark respiration of plants of

Correspondence: Mark G. Tjoelker, fax +1/979 8456049, e-mail m-tjoelker@tamu.edu

diverse biomes are also analysed and a general function that describes the short-term temperature dependence of Q_{10} for foliar respiration is presented.

Materials and methods

Five tree species common to the North American boreal forest were selected: quaking aspen (*Populus tremuloides* Michx.), paper birch (*Betula papyrifera* Marsh.), tamarack (*Larix laricina* [Du Roi] K. Koch), black spruce (*Picea mariana* [Mill.] B.S.P.), and jack pine (*Pinus banksiana* Lamb.). Plants were grown from seed in 2.7 dm³ pots in sand and fertilized daily with a complete nutrient solution. As part of a larger study, we grew the seedlings at three growth temperatures 18/12, 24/18, and 30/24 °C (16-h light/dark period) and two CO₂ concentrations of 580 and 370 μmol mol⁻¹. The CO₂ and temperature treatments were applied as a complete factorial in a set of six identical growth

chambers with high-intensity-discharge lighting, providing 1200 μmol m⁻² s⁻¹ PPFD (Conviron E15, Controlled Environments, Inc., Winnipeg, Manitoba, Canada). Plants of each species grew separately in 10–15 pots in each growth chamber and treatments lasted 97 days (Tjoelker *et al.* 1998a).

The objective of this study is to examine the Q_{10} of respiration in relation to measurement temperature. The larger study explored growth, photosynthesis, and respiration responses to the treatments (Tjoelker *et al.* 1998a,b; Tjoelker *et al.* 1999a,b). The analysis here is based on temperature-response functions for dark respiration determined for each species throughout the experiment (Tjoelker *et al.* 1999b). Specific rates of dark net CO₂ efflux were determined for fully developed foliage at measurement temperatures of 12, 18, 24 and 30 °C using portable infra-red gas analysers and cuvettes (LCA-3 and PLC-C, Analytical Development Co. Ltd, Hoddesdon, UK). The temperature responses for three replicate shoots, consisting of leaves or needles and attached stem, were determined for each species and treatment combination.

Because the determination of the temperature responses involved multiple measures on individual plants, we used a statistical analysis of variance that included both between and within plant variation. To this end we conducted a repeated-measures analysis of variance for each species. The full general linear model included the fixed effects of CO₂ (1 d.f.), temperature treatment (2 d.f.) and their interaction, and the random effect of shoot nested within treatment combination (12 d.f.). Measurement temperature was included in the model as a nested and fixed effect (3 d.f.) and partitioned into single-degree-of-freedom contrasts for linear and quadratic effects.

Inspection of model fits to natural-log transformed rates revealed a consistent departure from linearity in each species. Quadratic contrasts consistently showed the temperature response (on a log scale) to be a second-order polynomial (Fig. 1). In contrast to a linear fit, a second-order polynomial fit means that the slope (i.e. temperature coefficient) varies as a linear function of temperature. By definition, the temperature coefficient multiplied by 10 is equivalent to the natural log of the Q_{10} . Thus, the temperature coefficient or Q_{10} are dependent upon measurement temperature.

Given the widespread usage of Q_{10} in modelling and to facilitate comparisons across published studies, we show Q_{10} values to describe the temperature sensitivity of respiration. We calculated Q_{10} values separately for each of the three measurement temperature intervals (12–18, 18–24, and 24–30 °C) based on the equation:




Fig. 1 Short-term temperature response of foliar dark respiration for seedlings of five boreal tree species. Least squares means of natural log-transformed specific respiration rates are shown averaged across treatment combinations for each species (see Materials and methods). Quadratic polynomial fits for each species indicate a declining slope and Q_{10} with increasing measurement temperature. Standard errors of the least squares means are: *Populus tremuloides*, 0.024; *Betula papyrifera*, 0.022; *Larix laricina*, 0.026; *Pinus banksiana*, 0.022; and *Picea mariana*, 0.030.

$$R_{s(t)} = R_0 Q_{10}^{\left(\frac{t-t_0}{10}\right)}$$

where respiration rate, $R_{s(t)}$, at temperature t , is a function of R_0 , the specific respiration rate at a reference temperature t_0 (12, 18, or 24 °C), and Q_{10} , the ratio of rates given a 10 °C change in temperature. In order to equalize variances, the interval values of Q_{10} were transformed using the natural logarithm and the repeated-measures model was used to test the effect of measurement temperature.

Results and discussion

Decreasing slopes of natural log-transformed rates of respiration were found that were a function of increasing measurement temperature for each species (Fig. 1). The quadratic contrasts of the measurement temperature effect were statistically significant ($P \leq 0.002$). Therefore, for each of the five boreal tree species, calculated values of Q_{10} declined with increasing measurement-temperature intervals (Table 1). Similar declines in Q_{10} values were observed in plants in all temperature or CO₂ treatment combinations (Table 1). Moreover, the respiratory Q_{10} for five boreal tree species was independent of the temperature or CO₂ concentration at which the seedlings were grown (Tjoelker *et al.* 1999b).

Table 1 Temperature effect on calculated Q_{10} of dark respiration of five common tree species of the boreal forest biome grown at two CO₂ concentrations*

Species	Growth [CO ₂] [†]	Q_{10} of dark respiration Measurement temperature interval		
		12–18 °C	18–24 °C	24–30 °C
<i>Populus tremuloides</i>	370	2.26	1.95	1.62
	580	2.23	1.97	1.63
<i>Betula papyrifera</i>	370	2.48	2.27	2.01
	580	2.77	2.37	1.88
<i>Larix laricina</i>	370	2.68	2.34	1.81
	580	2.49	2.44	1.96
<i>Pinus banksiana</i>	370	2.48	2.37	1.88
	580	2.26	2.26	1.91
<i>Picea mariana</i>	370	2.61	2.18	1.93
	580	2.88	2.38	1.95

*Rates of dark net CO₂ efflux of foliage were determined at measurement temperatures of 12, 18, 24 and 30 °C to obtain short-term temperature responses and estimates of Q_{10} which differed among measurement temperature intervals in each species ($P \leq 0.017$).

[†]Growth CO₂ concentration ($\mu\text{mol mol}^{-1}$), growth temperature (18/12, 24/18, and 30/24 °C), and the interaction term had no effect on Q_{10} in any species ($P \geq 0.15$).

These findings were corroborated by a synthesis of published results of Q_{10} for foliar respiration rates consisting of 238 data points for 56 plant species (Table 2). The present literature data analysis included species of diverse taxa, including grasses, forbs, and woody plants of tropical, temperate, boreal, and arctic biomes. Measures included those of field- and laboratory-grown plants in a variety of thermal environments. Regardless of the thermal environment or species, the Q_{10} of respiration was dependent upon the range of measurement temperatures in a simple linear relationship (slope = -0.046 ± 0.0033 SEM) that was consistent among the biomes (Fig. 2). In addition, a dataset ($n = 32$) of respiration rates of roots was examined that revealed a comparable slope relating Q_{10} to measurement temperature (-0.042 ± 0.011 SEM), but lower intercept (2.85) compared to foliage (data not shown).

A declining Q_{10} indicates that the temperature sensitivity of respiration is reduced with increasing measurement temperature. For example, a 10 °C increase in measurement temperature at low measurement temperatures from 5 to 15 °C results in an approximate 2.8 fold increase in respiration rate, whereas an increase in temperature from 25 to 35 °C results in a less than two-fold (1.8) increase in rate. In practice, the use of a Q_{10} at 2.0 could lead to over-predictions of respiration with warming at higher temperatures (e.g. >27 °C, where $Q_{10} < 2.0$) and under-predicting increases in respiration with warming at lower measurement temperatures (e.g. <27 °C). Predicted temperature responses using a fixed vs. temperature-corrected Q_{10} will differ in magnitude and direction depending upon both the measurement and prediction temperatures. For example, using a fixed Q_{10} of 2.0 for a respiration rate determined at 10 °C results in predicted rates that are 17% larger at 5 °C and 15% lower at 15 °C compared to a model with a temperature-corrected Q_{10} of 2.8 for 10 °C. In the absence of a measured Q_{10} or if Q_{10} is known at a given temperature, it is suggested that the general linear relationship (or slope) may be used to select a temperature-corrected Q_{10} for the relevant temperature range for modelling. For example, the linear relationship of Q_{10} with measurement temperature (Fig. 2) could be substituted for the Q_{10} term in the temperature-response function. Consequently, climate data could be used to generate daily or seasonally adjusted Q_{10} values.

The temperature dependence of Q_{10} calls into question the widespread use of a constant Q_{10} in modelling rates of respiration and perhaps the accuracy of those predictions, including global estimates of terrestrial carbon balance. For example, a fixed Q_{10} of 2.0 is used in biogeochemical models such as Biome-BGC and Century (Schimel *et al.* 1997) as well as PnET (Aber & Federer 1992). In comparison, the TEM model does recognize a need to

Table 2 Literature survey of temperature coefficients (Q_{10}) of the short-term response of foliar dark respiration to measurement temperatures from 5 to 35°C

Species	Temperature class midpoint (°C)*							Reference and notes
	5	10	15	20	25	30	35	
Arctic								
<i>Antennaria alpina</i>	4.2	–	3.3	–	2.0	–	1.6	Wager (1941)
<i>Cassiope hypnoides</i>	2.1	–	2.2	–	2.2	–	2.1	
<i>Cassiope tetragona</i>	3.3	–	2.9	–	2.2	–	1.7	
<i>Diapensia lapponica</i>	3.8	–	3.4	–	2.1	–	2.1	
<i>Empetrum nigrum</i>	3.5	–	2.7	–	2.3	–	2.0	Lab
	3.6	–	2.7	–	2.2	–	–	Field
<i>Oxyria digyna</i>	3.1	–	1.6	–	–	–	–	
<i>Poa</i> spp.	2.7	–	2.3	–	–	–	–	
<i>Polygonum viviparum</i>	3.0	–	2.0	–	–	–	–	
<i>Ranunculus glacialis</i>	3.6	–	1.4	–	–	–	–	
<i>Ranunculus pygmaeus</i>	2.8	–	3.1	–	2.1	–	–	
<i>Salix herbacea</i>	3.5	–	2.5	–	–	–	–	
<i>Saxifraga cernua</i>	2.3	–	3.2	–	–	–	–	
<i>Saxifraga oppositifolia</i>	3.9	–	2.5	–	2.1	–	1.8	Lab
	3.0	–	2.7	–	2.2	–	1.9	Field
<i>Viscaria alpina</i>	2.8	–	2.2	–	2.0	–	2.0	
<i>Chamaenerium latifolium</i>	2.2	–	2.45	–	–	–	–	Müller (1928)‡
<i>Salix glauca</i>	3.17	–	1.79	–	–	–	–	
Boreal †								
<i>Betula papyrifera</i>	–	–	–	–	–	1.78	–	Ranney & Peet (1994)
<i>Picea abies</i>	–	2.5	2.0	–	–	–	–	Stockfors & Linder (1998)
Temperate								
<i>Betula nigra</i>	–	–	–	–	–	1.37	–	Ranney & Peet (1994)
<i>Chamaecyparis obtusa</i>	3.14	2.69	2.30	1.97	1.69	–	–	Paembonan <i>et al.</i> (1991)
<i>Eucalyptus pauciflora</i>	–	–	2.6	–	–	1.9	–	Kirschbaum & Farquhar (1984)
<i>Helianthus annuus</i>	–	–	–	–	2.4	–	1.7	Brown & Escombe (1925)#
	3.3	–	2.2	–	–	–	–	Kidd <i>et al.</i> (1921)#
<i>Lupinus</i> spp.	2.5	2.5	2.4	1.7	2.0	–	–	Clausen (1890)§
<i>Pisum sativum</i>	3.3	–	2.5	–	1.7	1.3	1.2	Fernandes (1923)#
	3	2.8	2.4	2.3	1.8	1.6	1.4	Kuijper (1910)¶
<i>Prunus laurocerasus</i>	–	–	2.2	–	2.3	–	–	Matthaei (1905)#
<i>Prunus persica</i>	–	–	–	–	2.11	–	2.03	Grossman & DeJong (1994)
<i>Solanum tuberosum</i>	3.3	2.1	1.9	2.2	2.1	1.8	2.1	Lundegårdh (1924)
<i>Triticum sativum</i>	2.9	2.4	2.1	1.9	1.6	1.2	1.1	Clausen (1890)¶
	3.5	–	–	–	1.5	–	–	Mayer (1876)#
	–	3.0	2.4	1.8	1.7	1.6	1.7	Rischavi (1876)#
	2.6	2.4	2.3	1.9	1.8	1.3	1.1	Kuijper (1910)¶
<i>Vicia faba</i>	–	–	2.5	–	2.1	–	–	Winter: Breeze & Elston (1978)
	–	–	1.7	–	1.5	–	–	Summer:
<i>Acer pennsylvanicum</i>	–	–	2.64	2.83	2.67	–	–	Bolstad <i>et al.</i> (1999), Mitchell <i>et al.</i> (1999) (and unpublished data)
<i>Acer rubrum</i>	–	–	2.83	2.82	2.63	–	–	
<i>Betula</i> spp.	–	–	3.57	2.98	2.62	–	–	
<i>Carya</i> spp.	–	–	2.75	2.59	2.40	–	–	
<i>Cornus florida</i>	–	–	3.05	3.07	2.60	–	–	
<i>Fraxinus</i> spp.	–	–	2.79	2.70	2.30	–	–	
<i>Liriodendron tulipifera</i>	–	–	2.58	2.30	2.10	–	–	
<i>Magnolia fraseri</i>	–	–	2.52	2.13	2.17	–	–	
<i>Nyssa sylvatica</i>	–	–	2.70	2.49	2.26	–	–	
<i>Oxydendron arboreum</i>	–	–	4.11	3.05	2.98	–	–	
<i>Platanus occidentalis</i>	–	–	2.46	2.38	2.11	–	–	
<i>Quercus alba</i>	–	–	2.36	2.30	2.17	–	–	
<i>Quercus coccinea</i>	–	–	2.37	2.49	2.40	–	–	

Table 2 Continued

Species	Temperature class midpoint (°C)*							Reference and notes
	5	10	15	20	25	30	35	
<i>Quercus prinus</i>	–	–	2.27	2.22	2.27	–	–	
<i>Quercus rubra</i>	–	–	2.41	2.48	2.37	–	–	
<i>Rhododendron maximum</i>	–	–	2.98	2.59	2.57	–	–	
<i>Robinia pseudoacacia</i>	–	–	2.85	2.67	2.50	–	–	
<i>Tilia americana</i>	–	–	2.55	2.46	2.27	–	–	
<i>Dactylus glomerata</i>	–	–	–	1.4	1.3	–	–	15 °C, growth temperature: Ziska & Bunce (1994)
	–	–	–	1.6	1.5	–	–	20
	–	–	–	1.9	1.7	–	–	25
	–	–	–	2.5	1.6	–	–	30
<i>Medicago sativa</i>	–	–	–	1.8	1.6	–	–	15
	–	–	–	1.8	1.6	–	–	20
	–	–	–	2.0	1.7	–	–	25
	–	–	–	2.2	1.8	–	–	30
Tropical								
<i>Calophyllum inophyllum</i>	–	–	1.8	–	1.9	–	1.9	Stocker (1935)
<i>Cassia fistula</i>	–	–	2.0	–	2.0	–	2.1	Shade leaf
	–	–	2.0	–	2.0	–	2.6	Sun leaf
<i>Stelechocarpus burahol</i>	–	–	3.0	–	2.0	–	1.7	Shade leaf
	–	–	2.7	–	2.8	–	1.8	Sun leaf
Mean	3.12	2.55	2.50	2.28	2.05	1.54	1.79	
Upper 95% C.I.	3.32	2.78	2.62	2.40	2.13	1.74	1.96	
Lower 95% C.I.	2.91	2.32	2.39	2.16	1.96	1.34	1.62	
N	27	8	65	43	65	9	21	

*Values are tabulated based on temperature class midpoints (measurement range typically 10 °C)

†The boreal data and summary statistics include the five species examined in this study (Table 1)

Based on data shown in: ‡Stocker 1935; #Wager 1941; ¶James 1953; §Forward 1960

adjust Q_{10} . In modelling the temperature dependence of respiration, TEM incorporates a linear decrease in Q_{10} from 2.5 to 2.0 between 0 and 5 °C, a constant value of 2.0 between 5 and 20 °C, and a linear decrease from 2.0 to 1.5 between 20 and 40 °C (McGuire *et al.* 1992). However, these values are lower overall and do not capture the linear response of the general relationship between temperature and respiratory Q_{10} . The consequences of using a fixed vs. temperature-dependent Q_{10} for integrated carbon budgets await testing and evaluation in these and other models.

In addition, it is suggested that care should be taken when comparing Q_{10} values across diverse thermal environments. In the present literature analysis, mean Q_{10} values across all temperatures were 2.14 for tropical, 2.26 for temperate, 2.20 for boreal, and 2.56 for arctic biomes. However, adjusting the means for the three biomes that shared a common covariate relationship with short-term measurement temperature ($P = 0.50$ for homogeneity of slopes test) revealed similar Q_{10} values at 19 °C

of 2.31 for temperate, 2.22 for boreal, and 2.42 for arctic biomes. An analysis of the limited tropical data ($n = 3$ species in one study) biome showed no relationship with temperature; however, more data are needed for a robust comparison with the other biomes. This analysis suggests that apparent biome differences in Q_{10} may, in part, be a result of measurement temperature differences. Studies at common temperatures are needed to distinguish the effect of measurement temperature from potential adaptive or acclimation responses of vegetation of contrasting climate zones. For example, the Q_{10} values of arctic and tropical lichens were comparable at common measurement temperatures (Scholander *et al.* 1952). In contrast, temperature coefficients of respiration were found to vary with climate of origin among woody species, but not among annuals or herbaceous perennials compared at common measurement temperatures (Criddle *et al.* 1994).

A temperature-dependent Q_{10} may complicate interpretation of annual variation in respiration in seasonal




Fig. 2 Q_{10} of respiration rates of plants as a function of measurement temperature. Symbols are the mean Q_{10} of species of arctic (16 species), boreal (6 species), temperate (31 species), and tropical (3 species) biomes plotted at the midpoints in 5 °C classes. Error bars indicate ± 1 SD of the class means of all observations. Data are of 56 species from 23 studies, including our experiment with five boreal tree species (Table 1,2). Based on covariance analysis, neither slopes ($P=0.50$) nor intercepts ($P=0.07$) of linear regressions differed among the arctic, boreal, and temperate biome groups, evidence for a common relationship across these biomes. The Q_{10} of the tropical species showed no relationship with temperature ($P=0.33$). A single linear regression was fit ($\pm 95\%$ C.I., dashed lines) to all data points ($R^2=0.45$, $n=238$, $P<0.0001$).

climates. A three-year record of aboveground night-time respiration of *Chamaecyparis obtusa* (Paembonan *et al.* 1991) revealed that Q_{10} varied as a function of annual variation in mean monthly night temperature, declining linearly with increasing ambient air temperatures. Although interpreted as temperature acclimation of Q_{10} (Paembonan *et al.* 1991; Larcher 1995), the linear response of long-term (seasonal) variation in Q_{10} with temperature was similar to the short-term response to measurement temperature shown in the present study. Furthermore, seasonal variation in Q_{10} based on measures at ambient temperatures alone is insufficient evidence of thermal acclimation, unless accompanied by measurements at standard temperatures. In comparison, the present authors' laboratory data of five boreal tree species suggest that Q_{10} may not acclimate to temperature (Tjoelker *et al.* 1999b). However, field studies of foliar

respiration of *Picea abies* (Stockfors & Linder 1998) and *Eucalyptus pauciflora* (Atkin *et al.* 2000b) each showed increased Q_{10} in autumn or winter than summer. These reports suggest that Q_{10} may acclimate to seasonal changes in temperature by increasing at lower temperatures in a pattern similar to that of the short-term temperature-response demonstrated here. It is apparent that both measurement-temperature and thermal environment together may influence respiratory Q_{10} . Further studies employing standard measurement temperatures are needed to discern daily or seasonal acclimation responses of respiratory metabolism from those of short-term temperature-responses (Atkin *et al.* 2000b).

Realized respiration rates are not only a function of short-term temperature responses, but respiration may also acclimate to thermal environment. Respiratory carbon exchange rates in plants are known to acclimate with time (hours to days) to prevailing temperatures (Larigauderie & Körner 1995; Tjoelker *et al.* 1999a,b; Atkin *et al.* 2000b). In effect, thermal acclimation to warmer environments results in a downward adjustment in respiration rates determined at a standard measurement temperature. Consequently, long-term responses of plant respiration to climate warming will differ from predictions based solely on short-term response functions (Atkin *et al.* 2000a,b). However, acclimation of the temperature-response function to thermal environment may occur largely through adjustment in the intercept (R_0) and less so through changes in the temperature coefficient or Q_{10} (Butler & Landsberg 1981; Tjoelker *et al.* 1999b; but see Atkin *et al.* 2000b). Although respiratory Q_{10s} (Table 2) and specific respiration rates differ markedly among plant species (Reich *et al.* 1998) and their growth environments, the short-term sensitivity of Q_{10} to measurement temperature appears to be a general phenomenon, apparently reflecting a consistent bias in the standard exponential Q_{10} model.

At the global scale, climatic controls appear important in limiting the productivity of biomes (Churkina & Running 1998). Biogeochemical models of combined effects of elevated atmospheric CO_2 concentrations and climate changes on terrestrial net primary productivity generally predict increases in production and carbon stocks (Cao & Woodward 1998; Pan *et al.* 1998). However, a positive feedback between temperature and the release of CO_2 to the atmosphere by terrestrial respiration could significantly reduce the magnitude of terrestrial carbon accumulation (Houghton *et al.* 1998). Respiration is a major determinant of the carbon balance of forests (Valentini *et al.* 2000). In boreal forest ecosystems, for example, annual carbon losses to autotrophic respiration were about 60% of the carbon fixed in photosynthesis (Ryan *et al.* 1997). Therefore, accurate modelling of respiration will be critical in estimating global carbon

balance. Given the major role of respiratory CO_2 efflux as a component of the net carbon balance of plants and ecosystems and its strong dependence on temperature, the use of a temperature-corrected Q_{10} as presented here may improve the accuracy of predictions of respiration from leaf to global scales.

Acknowledgements

This study was supported in part by a grant (IBN-9630241) from the National Science Foundation (USA). We thank Katherine Mitchell and Paul Bolstad for sharing unpublished data.

References

- Aber JD, Federer CA (1992) A generalized, lumped-parameter model of photosynthesis, evapotranspiration and net primary production in temperate and boreal forest ecosystems. *Oecologia*, **92**, 463–474.
- Amthor JS (1984) The role of maintenance respiration in plant growth. *Plant, Cell and Environment*, **7**, 561–569.
- Arrhenius S (1889) Über die Reaktionsgeschwindigkeit bei der Inversion von Rohrzucker durch Säuren. *Zeitschrift für Physikalische Chemie*, **4**, 226–248.
- Atkin OK, Edwards EJ, Loveys BR (2000a) Response of root respiration to changes in temperature and its relevance to global warming. *New Phytologist*, **147**, 141–154.
- Atkin OK, Holly C, Ball MC (2000b) Acclimation of snow gum (*Eucalyptus pauciflora*) leaf respiration to seasonal and diurnal variations in temperature: the importance of changes in the capacity and temperature sensitivity of respiration. *Plant, Cell and Environment*, **23**, 15–26.
- Belehrádek J (1930) Temperature coefficients in biology. *Biological Reviews*, **5**, 30–58.
- Berry JA, Raison JK (1981) Responses of macrophytes to temperature. In: *Encyclopedia of Plant Physiology, New Series*, (eds Lange OL *et al.*), Vol. 12A, pp. 277–338. Springer, Berlin.
- Bolstad PV, Mitchell K, Vose JM (1999) Foliar temperature-respiration response functions for broad-leaved tree species in the southern Appalachians. *Tree Physiology*, **19**, 871–878.
- Breeze V, Elston J (1978) Some effects of temperature and substrate content upon respiration and the carbon balance of field beans (*Vicia faba* L.). *Annals of Botany*, **42**, 863–876.
- Butler DR, Landsberg JJ (1981) Respiration rates of apple trees estimated by CO_2 -efflux measurements. *Plant, Cell and Environment*, **4**, 153–159.
- Cao MK, Woodward FI (1998) Dynamic responses of terrestrial ecosystem carbon cycling to global climate change. *Nature*, **393**, 249–252.
- Churkina G, Running SW (1998) Contrasting climatic controls on the estimated productivity of global terrestrial biomes. *Ecosystems*, **1**, 206–215.
- Collier DE, Cummins WR (1990) The effects of low growth and measurement temperature on the respiratory properties of five temperate species. *Annals of Botany*, **65**, 533–538.
- Cramer W, Kicklighter DW, Bondeau A *et al.* (1999) Comparing global models of terrestrial net primary productivity (NPP): overview and key results. *Global Change Biology*, **5**, 1–15 (Suppl.).
- Criddle RS, Hopkin MS, McArthur ED, Hansen LD (1994) Plant distribution and the temperature coefficient of metabolism. *Plant, Cell and Environment*, **17**, 233–243.
- Farrar JF, Williams ML (1991) The effects of increased atmospheric carbon dioxide and temperature on carbon partitioning, source-sink relations and respiration. *Plant, Cell and Environment*, **14**, 819–830.
- Forward DF (1960) Effect of temperature on respiration. In: *Handbuch der Pflanzenphysiologie* (eds Ruhland W *et al.*), Vol. 12, Part 2, pp. 234–258. Springer, Berlin.
- Grossman YL, DeJong TM (1994) Carbohydrate requirements for dark respiration by peach vegetative organs. *Tree Physiology*, **14**, 37–48.
- Houghton RA, Davidson EA, Woodwell GM (1998) Missing sinks, feedbacks, and understanding the role of terrestrial ecosystems in the global carbon balance. *Global Biogeochemical Cycles*, **12**, 25–34.
- James WO (1953) *Plant Respiration*. Clarendon Press, Oxford.
- Kirschbaum MUF, Farquhar GD (1984) Temperature dependence of whole-leaf photosynthesis in *Eucalyptus pauciflora* Sieb. ex Spreng. *Australian Journal of Plant Physiology*, **11**, 519–538.
- Larcher W (1995). *Physiological Plant Ecology: Ecophysiology and Stress Physiology of Functional Groups*, 3rd edn. Springer, Berlin.
- Larigauderie A, Körner C (1995) Acclimation of leaf dark respiration to temperature in alpine and lowland plant species. *Annals of Botany*, **76**, 245–252.
- Lloyd J, Taylor JA (1994) On the temperature dependence of soil respiration. *Functional Ecology*, **8**, 315–323.
- Lundegårdh H (1924) Der Temperaturfaktor bei Kohlensäureassimilation und Atmung. *Biochemische Zeitschrift*, **154**, 195–234.
- McGuire AD, Melillo JM, Joyce LA, Kicklighter DW, Grace AL, Moore B III, Vorosmarty CJ (1992) Interactions between carbon and nitrogen dynamics in estimating net primary productivity for potential vegetation in North America. *Global Biogeochemical Cycles*, **6**, 101–124.
- Melillo JM, McGuire AD, Kicklighter DW, Moore B III, Vorosmarty CJ, Schloss AL (1993) Global climate change and terrestrial net primary production. *Nature*, **363**, 234–240.
- Mitchell KA, Bolstad PV, Vose JM (1999) Interspecific and environmentally induced variation in foliar dark respiration among eighteen southeastern deciduous tree species. *Tree Physiology*, **19**, 861–870.
- Paembonan SA, Hagihara A, Hozumi K (1991) Long-term measurement of CO_2 release from the aboveground parts of a hinoki forest tree in relation to air temperature. *Tree Physiology*, **8**, 399–405.
- Pan Y, Melillo JM, McGuire AD *et al.* (1998) Modeled responses of terrestrial ecosystems to elevated atmospheric CO_2 : a comparison of simulations by the biogeochemistry models of the Vegetation/Ecosystem Modeling and Analysis Project (VEMAP). *Oecologia*, **114**, 389–404.
- Ranney TG, Peet MM (1994) Heat tolerance of five taxa of birch (*Betula*): Physiological responses to supraoptimal leaf temperatures. *Journal of the American Society of Horticultural Science*, **119**, 243–248.
- ap Rees T, Burrell MM, Entwistle TG, Hammond JBW, Kirk D, Kruger NJ (1988) Effects of low temperature on the respiratory metabolism of carbohydrates by plants. In: *Plants and Temperature* (eds Long SP, Woodward FI), Vol. 32, pp. 377–393. Society for Experimental Biology, Cambridge.

- Reich PB, Walters MB, Ellsworth DS, Vose JS, Volin JC, Gresham C, Bowman WD (1998) Relationships of leaf dark respiration to leaf nitrogen, specific leaf area and leaf life-span: a test across biomes and functional groups. *Oecologia*, **114**, 471–482.
- Ryan MG (1991) Effects of climate change on plant respiration. *Ecological Applications*, **1**, 157–167.
- Ryan MG, Lavigne MB, Gower ST (1997) Annual carbon cost of autotrophic respiration in boreal forest ecosystems in relation to species and climate. *Journal of Geophysical Research*, **102**, 28,871–28,883.
- Schimel DS, VEMAP Participants, Braswell BH (1997) Continental scale variability in ecosystem processes: models, data, and the role of disturbance. *Ecological Monographs*, **67**, 251–271.
- Scholander PF, Flagg W, Walters V, Irving L (1952) Respiration in some arctic and tropical lichens in relation to temperature. *American Journal of Botany*, **39**, 707–713.
- Stocker O (1935) Assimilation und Atmung westjavanischer Tropenbäume. *Planta*, **24**, 402–445.
- Stockfors J, Linder S (1998) The effect of nutrition on the seasonal course of needle respiration in Norway spruce stands. *Trees*, **12**, 130–138.
- Tjoelker MG, Oleksyn J, Reich PB (1998a) Temperature and ontogeny mediate growth response to elevated CO₂ in seedlings of five boreal tree species. *New Phytologist*, **140**, 197–210.
- Tjoelker MG, Oleksyn J, Reich PB (1998b) Seedlings of five boreal tree species differ in acclimation of net photosynthesis to elevated CO₂ and temperature. *Tree Physiology*, **18**, 715–726.
- Tjoelker MG, Oleksyn J, Reich PB (1999a) Acclimation of respiration to temperature and CO₂ in seedlings of boreal tree species in relation to plant size and relative growth rate. *Global Change Biology*, **5**, 679–691.
- Tjoelker MG, Reich PB, Oleksyn J (1999b) Changes in leaf nitrogen and carbohydrates underlie temperature and CO₂ acclimation of dark respiration in five boreal tree species. *Plant, Cell and Environment*, **22**, 767–778.
- Valentini R, Matteucci G, Dolman AJ *et al.* (2000) Respiration as the main determinant of carbon balance in European forests. *Nature*, **404**, 861–865.
- Van't Hoff JH (1896) *Studies in Chemical Dynamics*. F. Muller, Amsterdam, 286pp.
- Wager HG (1941) On the respiration and carbon assimilation rates of some arctic plants as related to temperature. *New Phytologist*, **40**, 1–19.
- Ziska LH, Bunce JA (1994) Direct and indirect inhibition of single leaf respiration by elevated CO₂ concentrations: interaction with temperature. *Physiologia Plantarum*, **90**, 130–138.